


***Self-Reported Diagnosis of Rheumatoid Arthritis in Adults
Stratified by Gender and Age***

Gender	Quartile of C8 PFOA	AGE GROUP	Diagnosis of Rheumatoid Arthritis	Population in Quartile	Num of Reported	Percent in Quartile
Female	1	30-50	No	3012	2924	97.08%
Female	1	30-50	Yes	3012	88	2.92%
Female	1	51-older	No	2738	2491	90.98%
Female	1	51-older	Yes	2738	247	9.02%
Female	2	30-50	No	3036	2951	97.20%
Female	2	30-50	Yes	3036	85	2.80%
Female	2	51-older	No	2749	2543	92.51%
Female	2	51-older	Yes	2749	206	7.49%
Female	3	30-50	No	3015	2925	97.01%
Female	3	30-50	Yes	3015	90	2.99%
Female	3	51-older	No	2744	2532	92.27%
Female	3	51-older	Yes	2744	212	7.73%
Female	4	30-50	No	3022	2941	97.32%
Female	4	30-50	Yes	3022	81	2.68%
Female	4	51-older	No	2739	2549	93.06%
Female	4	51-older	Yes	2739	190	6.94%
Male	1	30-50	No	2601	2551	98.08%
Male	1	30-50	Yes	2601	50	1.92%
Male	1	51-older	No	2668	2494	93.48%
Male	1	51-older	Yes	2668	174	6.52%
Male	2	30-50	No	2621	2573	98.17%
Male	2	30-50	Yes	2621	48	1.83%
Male	2	51-older	No	2649	2516	94.98%
Male	2	51-older	Yes	2649	133	5.02%
Male	3	30-50	No	2603	2566	98.58%
Male	3	30-50	Yes	2603	37	1.42%
Male	3	51-older	No	2662	2522	94.74%
Male	3	51-older	Yes	2662	140	5.26%
Male	4	30-50	No	2608	2571	98.58%
Male	4	30-50	Yes	2608	37	1.42%
Male	4	51-older	No	2659	2536	95.37%
Male	4	51-older	Yes	2659	123	4.63%

NOTE: All results reported here are based on information as reported by participants and were not independently verified. The association between self-reported Rheumatoid Arthritis and C8/PFOA is unadjusted for any potential confounders or covariates. Caution in interpretation is warranted.

Self- Reported Diagnosis of Rheumatoid Arthritis in Female Adults Stratified by Age



Self- Reported Diagnosis of Rheumatoid Arthritis in Male Adults Stratified by Age

