
Self-Reported Diagnosis of Multiple Sclerosis in Adults Stratified by Gender and Age

Gender	Quartile of C8 PFOA	AGE GROUP	Diagnosis of Multiple Sclerosis	Population in Quartile	Num of Reported	Percent in Quartile
Female	1	30-50	No	3012	3001	99.63%
Female	1	30-50	Yes	3012	11	0.37%
Female	1	51-older	No	2738	2726	99.56%
Female	1	51-older	Yes	2738	12	0.44%
Female	2	30-50	No	3036	3025	99.64%
Female	2	30-50	Yes	3036	11	0.36%
Female	2	51-older	No	2749	2738	99.60%
Female	2	51-older	Yes	2749	11	0.40%
Female	3	30-50	No	3015	3002	99.57%
Female	3	30-50	Yes	3015	13	0.43%
Female	3	51-older	No	2744	2736	99.71%
Female	3	51-older	Yes	2744	8	0.29%
Female	4	30-50	No	3022	3006	99.47%
Female	4	30-50	Yes	3022	16	0.53%
Female	4	51-older	No	2739	2721	99.34%
Female	4	51-older	Yes	2739	18	0.66%
Male	1	30-50	No	2601	2595	99.77%
Male	1	30-50	Yes	2601	6	0.23%
Male	1	51-older	No	2668	2661	99.74%
Male	1	51-older	Yes	2668	7	0.26%
Male	2	30-50	No	2621	2617	99.85%
Male	2	30-50	Yes	2621	4	0.15%
Male	2	51-older	No	2649	2641	99.70%
Male	2	51-older	Yes	2649	8	0.30%
Male	3	30-50	No	2603	2598	99.81%
Male	3	30-50	Yes	2603	5	0.19%
Male	3	51-older	No	2662	2659	99.89%
Male	3	51-older	Yes	2662	3	0.11%
Male	4	30-50	No	2608	2604	99.85%
Male	4	30-50	Yes	2608	4	0.15%
Male	4	51-older	No	2659	2656	99.89%
Male	4	51-older	Yes	2659	3	0.11%

NOTE:All results reported here are based on information as reported by participants and were not independently verified. The association between self-reported Multiple Sclerosis and C8/PFOA is unadjusted for any potential confounders or covariates. Caution in interpretation is warranted.

Self- Reported Diagnosis of Multiple Sclerosis in Female Adults Stratified by Age



Self- Reported Diagnosis of Multiple Sclerosis in Male Adults Stratified by Age

