SELF-REPORTED DIAGNOSIS OF AMYOTROPHIC LS STRATIFIED BY GENDER AND AGE GROUP

ACE					
AGE GROUP (years)	GENDER	SELF-REPORTED	POPULATION	# REPORTED	PERCENT
0-10	Female	No	2690	2690	100.00%
0-10	Female	Yes	2690	0	0.000%
0-10	Male	No	2703	2703	100.00%
0-10	Male	Yes	2703	0	0.000%
11-20	Female	No	4770	4770	100.00%
11-20	Female	Yes	4770	0	0.000%
11-20	Male	No	4961	4961	100.00%
11-20	Male	Yes	4961	0	0.000%
21-30	Female	No	5408	5408	100.00%
21-30	Female	Yes	5408	0	0.000%
21-30	Male	No	4599	4599	100.00%
21-30	Male	Yes	4599	0	0.000%
31-40	Female	No	5791	5791	100.00%
31-40	Female	Yes	5791	0	0.000%
31-40	Male	No	4880	4880	100.00%
31-40	Male	Yes	4880	0	0.000%
41-50	Female	No	6413	6413	100.00%
41-50	Female	Yes	6413	0	0.000%
41-50	Male	No	5674	5674	100.00%
41-50	Male	Yes	5674	0	0.000%
51-60	Female	No	5257	5257	100.00%
51-60	Female	Yes	5257	0	0.000%
51-60	Male	No	5073	5072	99.980%
51-60	Male	Yes	5073	1	0.020%
61-70	Female	No	3329	3328	99.970%
61-70	Female	Yes	3329	1	0.030%
61-70	Male	No	3405	3405	100.00%
61-70	Male	Yes	3405	0	0.000%
71-80	Female	No	1571	1571	100.00%
71-80	Female	Yes	1571	0	0.000%
71-80	Male	No	1597	1597	100.00%
71-80	Male	Yes	1597	0	0.000%
>80	Female	No	559	559	100.00%
>80	Female	Yes	559	0	0.000%
>80	Male	No	350	350	100.00%
>80	Male	Yes	350	0	0.000%

NOTE 1: Because of the very low self-reported prevalence of ALS in this population, graphical summaries are not included.

NOTE 2: In order to enroll in the C8 Health Project, potential participants needed to travel to one of the established community-based screening sites. It was beyond the scope of the C8 Health Project to recruit and screen participants living in group residential facilities, such as nursing homes. Therefore, as participants in the C8 Health Project did not include those individuals more at risk for having ALS (those living in nursing homes), the self-reported prevalence of ALS may be lower than expected in a population of this size.