Self-Reported Clinical Symptoms: Difficulty Swallowing


All Age and Gender Groups

<u>NOTE:</u> Symptoms were self reported by participants and were not independently verified with medical records or by a health care professional. Limited information about the characteristics of the self-reported symptoms, such as severity, duration, or history, was collected. Inferences about population health or associations based on these self-reported symptoms should be made cautiously.


SELF-REPORTED CLINICAL SYMPTOMS BY GENDER AND AGE GROUP: DIFFICULTY SWALLOWING

| Gender | | | | Difficulty Swallowing | | | | Total |
|--------|-----------|-------|-------------|-----------------------|-----------|--------|-------|--------|
| | | | | Frequently | Sometimes | Rarely | Never | |
| Female | Age Group | 0-10 | Count | 17 | 91 | 256 | 2326 | 2690 |
| | | | Percent (%) | .6% | 3.4% | 9.5% | 86.5% | 100.0% |
| | | 11-20 | Count | 42 | 257 | 731 | 3740 | 4770 |
| | | | Percent (%) | .9% | 5.4% | 15.3% | 78.4% | 100.0% |
| | | 21-30 | Count | 63 | 391 | 1026 | 3928 | 5408 |
| | | | Percent (%) | 1.2% | 7.2% | 19.0% | 72.6% | 100.0% |
| | | 31-40 | Count | 106 | 553 | 1103 | 4029 | 5791 |
| | | | Percent (%) | 1.8% | 9.5% | 19.0% | 69.6% | 100.0% |
| | | 41-50 | Count | 233 | 828 | 1296 | 4056 | 6413 |
| | | | Percent (%) | 3.6% | 12.9% | 20.2% | 63.2% | 100.0% |
| | | 51-60 | Count | 190 | 786 | 1023 | 3258 | 5257 |
| | | | Percent (%) | 3.6% | 15.0% | 19.5% | 62.0% | 100.0% |
| | | 61-70 | Count | 124 | 437 | 595 | 2173 | 3329 |
| | | | Percent (%) | 3.7% | 13.1% | 17.9% | 65.3% | 100.0% |
| | | 71-80 | Count | 73 | 212 | 223 | 1063 | 1571 |
| | | | Percent (%) | 4.6% | 13.5% | 14.2% | 67.7% | 100.0% |
| | | >80 | Count | 26 | 73 | 87 | 373 | 559 |
| | | | Percent (%) | 4.7% | 13.1% | 15.6% | 66.7% | 100.0% |
| | Total | | Count | 874 | 3628 | 6340 | 24946 | 35788 |
| | | | Percent (%) | 2.4% | 10.1% | 17.7% | 69.7% | 100.0% |
| Male | Age Group | 0-10 | Count | 15 | 112 | 276 | 2300 | 2703 |
| | | | Percent (%) | .6% | 4.1% | 10.2% | 85.1% | 100.0% |
| | | 11-20 | Count | 26 | 217 | 681 | 4037 | 4961 |
| | | | Percent (%) | .5% | 4.4% | 13.7% | 81.4% | 100.0% |
| | | 21-30 | Count | 41 | 284 | 859 | 3415 | 4599 |
| | | | Percent (%) | .9% | 6.2% | 18.7% | 74.3% | 100.0% |
| | | 31-40 | Count | 76 | 355 | 918 | 3531 | 4880 |
| | | | Percent (%) | 1.6% | 7.3% | 18.8% | 72.4% | 100.0% |
| | | 41-50 | Count | 115 | 470 | 1040 | 4049 | 5674 |
| | | | Percent (%) | 2.0% | 8.3% | 18.3% | 71.4% | 100.0% |
| | | 51-60 | Count | 117 | 507 | 912 | 3537 | 5073 |
| | | | Percent (%) | 2.3% | 10.0% | 18.0% | 69.7% | 100.0% |
| | | 61-70 | Count | 84 | 353 | 530 | 2438 | 3405 |
| | | | Percent (%) | 2.5% | 10.4% | 15.6% | 71.6% | 100.0% |
| | | 71-80 | Count | 45 | 154 | 240 | 1158 | 1597 |
| | | | Percent (%) | 2.8% | 9.6% | 15.0% | 72.5% | 100.0% |
| | | >80 | Count | 16 | 34 | 47 | 253 | 350 |
| | | | Percent (%) | 4.6% | 9.7% | 13.4% | 72.3% | 100.0% |
| | Total | | Count | 535 | 2486 | 5503 | 24718 | 33242 |
| | | | Percent (%) | 1.6% | 7.5% | 16.6% | 74.4% | 100.0% |

SELF-REPORTED DIFFICULTY SWALLOWING IN FEMALES BY AGE GROUP



SELF-REPORTED DIFFICULTY SWALLOWING IN FEMALES BY AGE GROUP



SELF-REPORTED DIFFICULTY SWALLOWING BY GENDER AND AGE GROUP

